arXiv:1302.4756v1 [math.AG] 19 Feb 2013

THE EXCEPTIONAL SET AND THE GREEN-GRIFFITHS LOCUS DO NOT ALWAYS COINCIDE

SIMONE DIVERIO AND ERWAN ROUSSEAU

Abstract. We give a very simple criterion in order to ensure that the Green-Griffiths locus of a projective manifold is the whole manifold. Next, we exhibit in all dimensions a projective hyperbolic manifold of general type, which satisfies such a criterion. In particular, this gives some precisions to an example given by M. Green to S. Lang.

1. introduction

Let X be a complex projective manifold, $V \subset T_X$ a rank r holomorphic subbundle, and call

$$E^{GG}_{k,m} V^* \to X$$

the holomorphic vector bundle of jet differentials of order k and weighted degree m acting on germs of holomorphic curves tangent to V. For definitions and basic properties of these bundles we refer to [GG80] and [Dem97]. Jet differentials are particularly useful in connection to hyperbolicity-related problems thanks to the following.

Theorem 1.1 (Green-Griffiths [GG80], Siu-Yeung [SY97], Demailly [Dem97]). *Let (X, V) be a directed projective manifold and $A \to X$ an ample line bundle. Then, for any entire curve $f\colon \mathbb{C} \to (X, V)$ tangent to V and any $P \in H^0(X, E^{GG}_{k,m} V^* \otimes A^{-1})$, one has $P(f', f'', \dots, f^{(k)}) \equiv 0$.*

Now, fix an ample line bundle $A \to X$. Define $GG_A(X, V)$ to be the set of points $x \in X$ such that for all integers $k > 0$ there exists a k-jet of holomorphic curve $\varphi_k\colon (\mathbb{C}, 0) \to (X, x)$ tangent to V with the property that for all integers $m > 0$, every global jet differential of order k and weighted degree m with values in A^{-1} vanishes whenever evaluated on $\varphi_k(0)$. Thus, if $f : \mathbb{C} \to X$ is an entire curve tangent to V, Theorem 1.1 tells us immediately that $f(\mathbb{C}) \subset GG_A(X, V)$. We shall see in the next section that the locus $GG_A(X, V)$ is indeed independent of the particular ample line bundle chosen.

Definition 1.2. Let X be a projective manifold and $V \subset T_X$ a holomorphic subbundle. The *exceptional set* $\mathrm{Exc}(X, V)$ of (X, V) is defined to be the Zariski closure of the union of all the images of entire curves traced in X and tangent to V. The *Green-Griffiths locus* $GG(X, V)$ *of* (X, V) is defined as

$$GG(X, V) = GG_A(X, V),$$

for some (and hence any) ample line bundle $A \to X$.

In the *absolute case* $V = T_X$, we shall simply call $\mathrm{Exc}(X, T_X) = \mathrm{Exc}(X)$ and $GG(X, T_X) = GG(X)$.

Date: February 21, 2013.

2010 *Mathematics Subject Classification.* Primary: 32Q45, 32M15; Secondary: 37F75.

Both authors are partially supported by the ANR project “POSITIVE”, ANR-2010-BLAN-0119-01.

By Theorem 1.1, one always has the inclusion:

$$\operatorname{Exc}(X,V)\subset GG(X,V).$$

It is conjectured in [GG80] that if X is a n dimensional projective manifold of general type and $V=T_X$, then there should exists a large integer $k=k(X)$ such that the growth of

$$m\mapsto \dim H^0(X,E^{GG}_{k,m}T^*_X)$$

would have to be maximal, that is asymptotic with $m^{(k+1)n-1}$. This conjecture was proven in [GG80] for projective surfaces and therein supported in all dimension by an Euler characteristic computation. It was then established in the special case of smooth projective hypersurfaces of general type in [Mer10] and was finally proven in full generality only very recently by J.-P. Demailly in [Dem11] by means of his holomorphic Morse inequalities combined with a delicate "probabilistic" curvature estimate. In particular, if X is a projective manifold of general type and $A\to X$ an ample line bundle, then there always exists large integers $k,m>0$ such that $H^0(X,E^{GG}_{k,m}T^*_X\otimes A^{-1})\neq\{0\}$ and indeed this space is very big. It is thus legitimate to ask if much more is true, namely that if X is a projective manifold of general type, then the Green-Griffiths locus is always a proper subvariety of X, *i.e.*

$$GG(X)\subsetneq X.$$

An affirmative answer to this would lead directly to the solution of the so-called Green-Griffiths-Lang conjecture. One could even speculate more and ask whether Exc(X) $=GG(X)$. About this, let us quote S. Lang from his paper [Lan86], page 200:

"In particular, the exceptional set [...] is contained in the Green-Griffiths set [...]. I asked Green-Griffiths whether they might be equal. Green told me that the two sets are not equal in general. Certain Hilbert modular surfaces constructed by Shavel [Sha78], compact quotients of the product of the upper half-plane with itself, provide a counterexample which is hyperbolic, but such that the Green-Griffiths set is the whole variety. Thus the jet construction appears insufficient so far to characterize the exceptional set [...] completely algebraically."

In [Lan86] there are no more details about that, nor in the subsequent literature as far as we know. Here, we take the opportunity to give a detailed account of the above fact by proving indeed a quite more general result.

Let $\mathcal{W}$ be a coherent analytic sheaf on a complex manifold X and $m>0$ be an integer. We shall denote by $\mathcal{W}^{[m]}$ the double dual of the m-th tensor power $\mathcal{W}^{\otimes m}$ of $\mathcal{W}$.

Theorem 1.3. *Let (X,V) be a complex projective directed manifold. Suppose that there exists a saturated coherent analytic subsheaf $\mathcal{W}\subset\mathcal{O}_X(V)$ such that for all ample line bundle $A\to X$ one has*

$$H^0(X,(\mathcal{W}^*)^{[m]}\otimes A^{-1})=0,\quad \forall m\geq 1.$$

Then, $GG(X,V)=X$.

Observe that we do not require any Frobenius integrability property for $\mathcal{W}$. Now, since a saturated subsheaf of a locally free sheaf is reflexive and since a reflexive rank one sheaf on a smooth complex manifold is invertible, in the special case of rank one subsheaves we have the following.

Corollary 1.4. *Let (X,V) be a complex projective directed manifold endowed with a rank one saturated coherent analytic subsheaf $\mathcal{O}_X(L)\subset\mathcal{O}_X(V)$, where $L\to X$ is a holomorphic line bundle. Suppose that L^* is not big. Then, $GG(X,V)=X$.*

This last corollary can be rephrased by saying that if there exists on X a holomorphic (possibly singular) foliation $\mathcal{F}$ by curves tangent to V whose canonical bundle $K_{\mathcal{F}} \simeq L^*$ is not big, then $GG(X,V) = X$.

Proof. It is an immediate consequence of Kodaira's lemma: a line bundle $F \to X$ is big if and only if for all ample line bundle $A \to X$ there exists an integer $k > 0$ such that

$$H^0(X, F^{\otimes k} \otimes A^{-1}) \neq \{0\}.$$

Thus, since $(\mathcal{O}_X(L)^*)^{[m]} \simeq \mathcal{O}_X(L^*)^{\otimes m}$, if L^* is not big then $\mathcal{O}_X(L)$ satisfies the requirements of Theorem 1.3 and we are done. ☐

Using Corollary 1.4 we obtain:

Theorem 1.5. *Let X be a compact free quotient of the n-dimensional polydisc by a subgroup of* $\mathrm{Aut}(\Delta)^n$. *Then,*

- *X is of general type*
- *the exceptional locus* $\mathrm{Exc}(X)$ *is empty*
- *the Green-Griffiths locus $GG(X)$ is the whole manifold.*

In the crucial case of surfaces, a similar kind of conclusions was already known to M. McQuillan from a completely foliated point of view, as he kindly pointed out to us. For precise statements and considerations we refer the reader to the web page [McQ12] and, in particular, to Items 10, 12 and 13 on that page.

Coming back to Theorem 1.5, the first two points of are immediate thanks to the following remarks.

Remark 1.6 (For details, see [Mok89]). Since Δ^n is a bounded homogeneous domain, we have that its Bergman metric is $\mathrm{Aut}(\Delta^n)$-invariant and also Kähler-Einstein (with negative Ricci curvature). In particular, it descends to a Kähler-Einstein metric with negative Ricci curvature on every smooth quotient X of Δ^n. Thus, the canonical bundle K_X of X is positive and in particular X is projective and of general type.

Remark 1.7. If X is a complex space whose universal cover is a bounded domain in $\mathbb{C}^n$, then every entire curve must be constant by Liouville's theorem. Hence, the exceptional locus Exc(X) is empty.

The third point will be discussed in Section 3. As a consequence of Theorem 1.5, we recover Green's counterexample and show as well that in order to get the same conclusion it is indeed sufficient to consider any compact free quotient of the n-dimensional polydisc by a subgroup acting diagonally. For instance, a product of compact hyperbolic Riemann surfaces will do.

Acknowledgments. Both authors are glad to thank Michael McQuillan for several interesting comments and explanations.

The first-named author friendly thanks Sébastien Boucksom and Nicolas Bergeron for several interesting and fruitful discussions and remarks, and J.-P. Demailly for suggesting to add Lemma 2.2.

The second-named author warmly thanks Steven Lu for very interesting discussions and remarks that greatly improved a previous version of this paper and for the indication of the reference [Lu96]. He also thanks the CNRS for the opportunity to spend a semester in Montreal at the UMI CNRS-CRM and the hospitality of UQAM-Cirget where part of this work was done.

2. Proof of Theorem 1.3

Let us start by recalling some basic facts from [Dem97] about jet differentials in the general framework of directed manifolds (for more details see the cited references).

So, let (X, V) be a directed manifold, that is a complex manifold X together with a holomorphic subbundle $V \subset T_X$, non necessarily integrable, of the holomorphic tangent bundle T_X of X. We call J_kV the bundle of k-jets of holomorphic curves $\varphi\colon (\mathbb{C}, 0) \to X$ which are tangent to V, that is $\varphi'(t) \in V_{\varphi(t)}$ for all t, together with the projection $\varphi \mapsto \varphi(0)$ onto X. It is a holomorphic fiber bundle, which is naturally a subbundle of the bundle J_kT_X of k-jets of holomorphic curves with values in X with no further restrictions. Moreover, there is a canonically defined fiber-wise $\mathbb{C}^*$-action on J_kV given by the reparametrization of a k-jet tangent to V by the homotheties corresponding to elements of $\mathbb{C}^*$. Of course, this action is compatible with restriction to smaller subbundles.

Let $E^{GG}_{k,m}V^* \to X$ be the holomorphic vector bundle over X whose fibers are complex valued polynomials on the fibers of J_kV of weighted degree m with respect to the above-defined $\mathbb{C}^*$-action. If $W \subset V \subset T_X$ are holomorphic subbundles, the inclusions

$$J_kW \subset J_kV \subset J_kT_X$$

induce surjective arrows

$$E^{GG}_{k,m}T^*_X \to E^{GG}_{k,m}V^* \to E^{GG}_{k,m}W^*,$$

so that $E^{GG}_{k,m}W^*$ can be regarded as a quotient of $E^{GG}_{k,m}V^*$, the quotient map being given by evaluating jet differentials tangent to V only on k-jets tangent to W.

Remark 2.1. In particular, we see that if every global section of $E^{GG}_{k,m}W^*$ vanishes at some point $j_k \in J_kW \subset J_kV$, then so every global section of $E^{GG}_{k,m}V^*$ does.

Now, considering the highest monomials (with respect to the reverse lexicographic order) gives a natural filtration on weighted homogeneous polynomials. Such filtration defines an intrinsic filtration on the bundles $E^{GG}_{k,m}V^*$, whose graded series are given by

$$\operatorname{Gr}^\bullet E^{GG}_{k,m}V^* = \bigoplus_{\ell_1+2\ell_2+\cdots+k\ell_k=m} S^{\ell_1}V^* \otimes \cdots \otimes S^{\ell_k}V^*.$$

Therefore, it follows that if for all k-tuple of non negative integers $(\ell_1, \dots, \ell_k)$ such that $\ell_1 + 2\ell_2 + \cdots + k\ell_k = m$, we have

$$H^0(X, S^{\ell_1}V^* \otimes \cdots \otimes S^{\ell_k}V^*) = 0,$$

then

$$H^0(X, E^{GG}_{k,m}V^*) = 0,$$

as well.

Before entering into the proof of Theorem 1.3, let us show as promised that the locus $GG_A(X, V)$ is independent of the ample line bundle $A \to X$.

Lemma 2.2. *The set $GG_A(X, V)$ does not depend on the ample line bundle $A \to X$.*

Proof. Let $A, B \to X$ two ample line bundles and $\ell > 0$ be a positive integer such that $B^{\otimes \ell} \otimes A^{-1}$ is globally generated. We shall show that

$$GG_A(X, V) \subset GG_B(X, V),$$

and the equality will follow by interchanging the roles of A and B.

Let $x \notin GG_B(X, V)$. Then, there exist integers $k, m > 0$, a global section $P \in H^0(X, E^{GG}_{k,m}V^* \otimes B^{-1})$ and a k-jet of holomorphic curve $\varphi_k\colon (\mathbb{C}, 0) \to (X, x)$

tangent to V such that $P(\varphi_k)\neq 0$. Let $\sigma\in H^0(X,B^{\otimes\ell}\otimes A^{-1})$ be a global section such that $\sigma(x)\neq 0$. Then,

$$P^\ell\otimes\sigma\in H^0(X,E^{GG}_{k,\ell m}V^*\otimes A^{-1})$$

and

$$(P^\ell\otimes\sigma)(\varphi_k)=P^\ell(\varphi_k)\cdot\sigma(x)\neq 0,$$

so that $x\notin GG_A(X,V)$. $\square$

Proof of Theorem 1.3. Let us first suppose that $\mathcal{W}=\mathcal{O}_X(W)$ is the sheaf of germs of holomorphic sections of a holomorphic subbundle $W\subset V$. Let $A\to X$ be any ample line bundle and $k,m>0$ be two integers. For all k-tuple of non-negative integers $(\ell_1,\dots,\ell_k)$ such that $\ell_1+2\ell_2+\cdots+k\ell_k=m$ we have

$$S^{\ell_1}W^*\otimes\cdots\otimes S^{\ell_k}W^*\otimes A^{-1}\subset(W^*)^{\otimes|\ell|}\otimes A^{-1},$$

where $|\ell|=\sum\ell_j$. Thus,

$$H^0(X,S^{\ell_1}W^*\otimes\cdots\otimes S^{\ell_k}W^*\otimes A^{-1})=0$$

and, since taking the tensor product of $E_{k,m}W^*$ with A^{-1} affects the corresponding graded bundle just tensoring by A^{-1}, we have

$$H^0(X,E^{GG}_{k,m}W^*\otimes A^{-1})=0.$$

Moreover, we have of course surjective morphisms

$$E^{GG}_{k,m}V^*\otimes A^{-1}\to E^{GG}_{k,m}W^*\otimes A^{-1}.$$

Therefore, by Remark 2.1, in such a situation every global jet differential tangent to V of order k and weighted degree m with values in A^{-1} must vanish when evaluated on k-jets which are tangent to W. Since at each point $x\in X$ we have such jets, it follows that $GG(X,V)=X$.

Now, take $\mathcal{W}$ as in the hypotheses. Since $\mathcal{W}$ is saturated, $\mathcal{O}_X(V)/\mathcal{W}$ is torsion-free and thus locally free in codimension two. In other words, there exists a proper subvariety $Y\subsetneq X$, $\operatorname{codim}_X Y\geq 2$, and a holomorphic vector bundle

$$W\to U:=X\setminus Y$$

on the dense open subset U such that $\mathcal{O}_U(W)\simeq\mathcal{W}|_U$ and W is actually a subbundle of the restriction $V|_U$ of V to U. Of course, over U, we also have $\mathcal{W}^*|_U\simeq\mathcal{O}_U(W^*)$. Since $(\mathcal{W}^*)^{[|\ell|]}$ is reflexive by definition and thus normal, we have a surjection

$$H^0(X,(\mathcal{W}^*)^{[|\ell|]}\otimes A^{-1})\to H^0(U,(\mathcal{W}^*)^{[|\ell|]}\otimes A^{-1})=H^0(U,(W^*)^{\otimes|\ell|}\otimes A^{-1}),$$

so that $H^0(U,(W^*)^{\otimes|\ell|}\otimes A^{-1})=\{0\}$. From

$$S^{\ell_1}W^*\otimes\cdots\otimes S^{\ell_k}W^*\otimes A^{-1}|_U\subset(W^*)^{\otimes|\ell|}\otimes A^{-1}|_U,$$

we conclude that

$$H^0(U,\operatorname{Gr}^\bullet E^{GG}_{k,m}W^*\otimes A^{-1})=\{0\}$$

and thus $H^0(U,E^{GG}_{k,m}W^*\otimes A^{-1})=\{0\}$. Then, since each section of $H^0(U,E^{GG}_{k,m}V^*\otimes A^{-1})$ extends to a section of $H^0(X,E^{GG}_{k,m}V^*\otimes A^{-1})$, we have $U\subset GG(X,V)$ and since $GG(X,V)$ is a closed set, then $GG(X,V)=X$. $\square$

3. Compact quotients of the polydisc: proof of Thoerem 1.5

Let X be a complex projective manifold as in the statement of Theorem 1.5. Thus, $X = \Delta^n/\Gamma$, where

$$\pi_1(X) = \Gamma \subset \mathrm{PSL}(2,\mathbb{R})^n = \underbrace{\mathrm{Aut}(\Delta) \times \cdots \times \mathrm{Aut}(\Delta)}_{n \text{ times}}$$

is co-compact and acts freely and properly discontinuously on Δ^n. By the so-called Density Lemma [Rag72, Cor.(5.21) and Thm.(5.22), p.86] the following dichotomy holds: either at least one among the n projections

$$\mathrm{pr}_i\colon \Gamma \to \mathrm{PSL}(2,\mathbb{R}), \quad i = 1,\dots,n,$$

of Γ to the i-th factor of $\mathrm{PSL}(2,\mathbb{R})^n$ has dense image, or X is up to finite Galois covers a product of compact hyperbolic Riemann surfaces, *i.e.* there is a normal subgroup $\Gamma_0 \trianglelefteq \Gamma$ of finite index in Γ such that Δ^n/Γ_0 is a product of compact hyperbolic Riemann surfaces. We first treat the latter case.

Since Γ as well as Γ_0 act diagonally, after possibly a reordering of the factors, the situation is the following

$$\begin{tikzcd}
\Delta^n \arrow[d, "\pi_1"] \arrow[dd, bend right=60, "\pi"'] & T_{\Delta^n} \simeq \bigoplus_{i=1}^n p_i^* T_\Delta \arrow[l] \\
C_1 \times \cdots \times C_n \arrow[d, "\pi_2"] & T_{C_1\times\cdots\times C_n} \simeq \bigoplus_{i=1}^n q_i^* T_{C_i} \arrow[l] \\
X & T_X \simeq \bigoplus_{i=1}^n L_i, \arrow[l]
\end{tikzcd}$$

where $p_i\colon \Delta^n \to \Delta$ and $q_i\colon C_1 \times \cdots \times C_n \to C_i$, $i = 1,\dots,n$, are the projections onto the i-th factor, and we have compatible isomorphisms

$$\pi_2^* L_i \simeq q_i^* T_{C_i}, \quad \pi^* L_i \simeq \pi_1^* \, q_i^* T_{C_i} \simeq p_i^* T_\Delta, \quad i = 1,\dots,n.$$

Since π_2 is a finite étale cover, the Kodaira-Iitaka dimension $\kappa(L_i^*)$ is the same of $\kappa(\pi_2^* L_i^*) = \kappa(q_i^* T_{C_i}^*) = 1$. In particular, any of the L_i provides a holomorphic line subbundle of T_X whose dual is not big, as soon as $n \geq 2$. Corollary 1.4 is thus verified in this case.

Next, we fix once and for all a co-compact subgroup $\Gamma \subset \mathrm{PSL}(2,\mathbb{R})^n$ acting freely and properly discontinuously on Δ^n, and such that the projection

$$\Gamma \to \mathrm{PSL}(2,\mathbb{R})$$

onto, say, the first factor has dense image Γ_1. Call $z \in \Delta$ the complex coordinate of the first factor and $w = (w_1,\dots,w_{n-1}) \in \Delta^{n-1}$ the complex coordinates of the last $n-1$ factors. We begin with a preliminary lemma.

Lemma 3.1 (Compare with [SB95])**.** *Let $\eta = f(z,w)\,(dz)^{\otimes m}$ be a symmetric differential of degree m on $\Delta_z \times \Delta_w^{n-1}$. Suppose that η is Γ-invariant. Then, η vanishes identically.*

This lemma should be regarded as a consequence of the following classical result for automorphic forms. Let $\Gamma \subset \mathrm{PSL}(2,\mathbb{R})^n$ be an irreducible discrete subgroup with compact quotient Δ^n/Γ, and f a Γ-automorphic form of weight $2r$, $r = (r_1,\dots,r_n) \in \mathbb{Z}^n$. If $r_1 \cdots r_n = 0$, then f must vanish identically (for this and related statements see for instance [Fre90]). Again by the Density Lemma, if for instance Γ is irreducible, then the image of Γ under any projection onto the factors will have dense image, so that any such irreducible Γ satisfies our hypotheses.

We shall come back about the existence of groups Γ as above in a moment.

Proof. Consider the smooth real function given by taking the Poincaré norm of η:

$$(z,w)\mapsto |f(z,w)|(1-|z|^2)^m.$$

This is a Γ-invariant smooth function, thus defined on the compact quotient $X=\Delta^n/\Gamma$. Let $p\in X$ be a point where this function attains its maximum and consider the discrete set of points $\{(z_{(j)},w_{(j)})\}_{j\in J}\in\Delta\times\Delta^{n-1}$ which are in the preimage of p by the quotient map. For each $j\in J$, the holomorphic map defined on the polydisc $\{z_{(j)}\}\times\Delta_w$ by

$$w\mapsto f(z_{(j)},w)$$

attains a maximum at the interior point $w_{(j)}$ and so it is constant. So, all the w_λ-derivatives vanish:

$$\partial f/\partial w_\lambda(z_{(j)},w)=0,\quad \text{for all } j\in J,\ \lambda=1,\dots,n-1, \text{ and } w\in\Delta^{n-1}.$$

By the density of Γ_1, the set $\{z_{(j)}\}_{j\in J}$ is dense in Δ, so $\partial f/\partial w_\lambda\equiv 0$, $\lambda=1,\dots,n-1$, and f does not depend on w. Therefore η does only depend on z and can thus be regarded as a symmetric differential of degree m on Δ_z, invariant by the action of Γ_1 on $\mathrm{Aut}(\Delta_z)$ which has, once again by hypothesis, dense image. But then, $\eta\equiv 0$. ☐

Now, consider the projective n-dimensional complex manifold $X=\Delta^n/\Gamma$: it is canonically polarized and Kobayashi-hyperbolic.

The holomorphic foliation by discs $\widetilde{\mathcal{F}}$ on Δ^n generated by $\partial/\partial z$ descends to a smooth foliation by curves $\mathcal{F}$ on X. Consider its tangent bundle $T_{\mathcal{F}}$: it is a rank one holomorphic subbundle of the tangent bundle T_X of X. The tangent bundle of the foliation will play the role of our line bundle L in the statement of the Corollary 1.4. In particular, we will see in a moment that its dual, the canonical bundle $K_{\mathcal{F}}=T^*_{\mathcal{F}}$ of $\mathcal{F}$ has negative Kodaira-Iitaka dimension.

Proposition 3.2 (Compare for instance with [Bru04])**.** *The canonical bundle $K_{\mathcal{F}}$ of $\mathcal{F}$ has negative Kodaira-Iitaka dimension.*

Proof. We have to show that $H^0(X,K_{\mathcal{F}}^{\otimes\ell})=\{0\}$ for all integers $\ell>0$. The latter space of global sections identifies canonically with the space of Γ-invariant global holomorphic sections of $K_{\widetilde{\mathcal{F}}}^{\otimes\ell}$ over Δ^n. Since $\widetilde{\mathcal{F}}$ is generated by $\partial/\partial z$, these sections are exactly of the form considered in Lemma 3.1, and therefore they vanish identically. ☐

Now, let us spend a few words about the existence of co-compact subgroups $\Gamma\subset\mathrm{PSL}(2,\mathbb{R})^n$ acting freely and properly discontinuously on Δ^n with dense projection onto the first factor, which is however quite classical.

First of all, observe that it is enough to consider the case $n=2$. Indeed, once such a subgroup Γ' of $\mathrm{PSL}(2,\mathbb{R})^2$ has been constructed, then we can take Γ to be

$$\Gamma=\Gamma'\times\pi_1(C_1)\times\cdots\times\pi_1(C_{n-2}),$$

where $\pi_1(C_j)$, $j=1,\dots,n-2$, is the fundamental group of a compact Riemann surface of genus $g(C_j)\geq 2$. The resulting quotient manifold will be then

$$S\times C_1\times\cdots\times C_{n-2},\quad S=\Delta^2/\Gamma',$$

and the image Γ_1 of the projection of Γ onto the first factor equals the projection Γ_1' of Γ', which is dense. Of course, less trivial examples (*i.e.* which do not split as a product even after finite étale covers) can be constructed as soon as one disposes of an irreducible co-compact subgroup of $\mathrm{PSL}(2,\mathbb{R})^n$ acting freely and properly discontinuously on Δ^n. Coming back to the surface case, such a Γ' is constructed for instance in [Sha78] (see also [Tak90]), and we give here a sketchy account for the sake of completeness.

3.1. **Construction of Γ'.** Take a quaternion algebra A which is division and whose center is a totally real quadratic number field k (for an excellent reference about quaternion algebras and Fuchsian groups see [Kat92]). Assume that

$$A \otimes_{\mathbb{Q}} \mathbb{R} = M(2, \mathbb{R})^2,$$

that is, A is unramified at the two places corresponding to the two different embeddings of k into $\mathbb{R}$. If $\mathfrak{O}$ is a maximal order in A, denote by $\Gamma(1)$ the group of units in $\mathfrak{O}$ with reduced norm 1, and identify it with its isomorphic image in $\mathrm{SL}(2, \mathbb{R})^2$. Next, call $\Gamma'' = \Gamma(1)/\{\pm 1\}$ the image of $\Gamma(1)$ in $\mathrm{PSL}(2, \mathbb{R})^2$. Then, it is well known that the action of Γ'' on Δ^2 is irreducible and properly discontinuous (see [Sha78, Shi94]); moreover, the fact that A is division classically implies that this action is co-compact.

Now, since the action is co-compact, Γ'' is a group of finite type. By Selberg's theorem, a finitely generated linear group over a field of zero characteristic is virtually torsion-free, *i.e.* it has some finite index subgroup Γ' which is torsion-free. Thus, Γ' acts freely on Δ^2. Moreover, being of finite index in Γ'', it is straightforward to see that it is again irreducible and with a properly discontinuous, co-compact action.

3.2. **Surfaces of general type with a holomorphic foliation by curves.** Let S be a surface of general type and $L \subset T_S$ a holomorphic line subbundle, *i.e.* S is endowed with a smooth holomorphic foliation $\mathcal{F}$ by curves whose tangent bundle $T_{\mathcal{F}}$ is L. Suppose, as in Corollary 1.4, that the canonical bundle $K_{\mathcal{F}}$ is not big.

Proposition 3.3. *Let $(S, \mathcal{F})$ be as above. Then, S is a quotient of the bidisc Δ^2.*

Proof. Following [Bru97], the existence of such a smooth foliation on the surface of general type S implies that K_S is ample and so, by Aubin-Yau, S is Kähler-Einstein and hence T_S is K_S-semistable. The semistability inequality reads

$$c_1(T_{\mathcal{F}}) \cdot c_1(S) \geq \frac{1}{2}\, c_1(S)^2 > 0. \tag{1}$$

Since $\mathcal{F}$ is smooth, the Baum-Bott formulae give

$$\begin{aligned} &c_2(S) - c_1(T_{\mathcal{F}}) \cdot c_1(S) + c_1(T_{\mathcal{F}})^2 = 0 \\ &c_1(S)^2 - 2\, c_1(T_{\mathcal{F}}) \cdot c_1(S) + c_1(T_{\mathcal{F}})^2 = 0. \end{aligned} \tag{2}$$

If T_S is stable, the first inequality in (1) is strict and using the second of (2) we obtain $c_1(T_{\mathcal{F}})^2 > 0$. Thus, $T_{\mathcal{F}}$ or its dual $K_{\mathcal{F}}$ must be big. Since, by (1), $c_1(K_{\mathcal{F}}) \cdot c_1(K_S) > 0$, we get that $K_{\mathcal{F}}$ is big, contradiction.

Therefore, T_S is polystable but not stable, that is $T_S = L_1 \oplus L_2$ is a direct sum of two line bundles and by Beauville-Yau's uniformization theorem [Bea00, Yau93], the universal cover $\widetilde{S}$ of S splits as a product of simply connected Riemann surfaces, the decomposition of the tangent bundle lifts and the fundamental group of S acts diagonally on $\widetilde{S}$. Of course, the only possibility for $\widetilde{S}$ is to be the product of two discs. □

It is known by [Lu96] that the surfaces as above satisfy the Green-Griffiths conjecture since they verify the Chern numbers inequality $c_1(S)^2 - 2\, c_2(S) \geq 0$ (to see this just inject the difference of the two identities in (2) into the first inequality in (1)).

Next, let us put ourself in the most general case covered by Corollary 1.4 in the case of surfaces of general type. So, assume that S is a smooth projective surface of general type and $\mathcal{F}$ a (possibly singular, with at most isolated singularities) holomorphic foliation by curves whose canonical bundle $K_{\mathcal{F}}$ is not big. By Saidenberg's theorem, we can suppose without loss of generality that $\mathcal{F}$ has reduced singularities.

Thus, the birational classification of foliations developed by Brunella and McQuillan (see [Bru97, McQ08] and [Bru04]) tells us that $\mathcal{F}$ is necessarily of the following two types.

- A Hilbert modular foliation, and thus S is a Hilbert modular surface, if $\kappa(K_{\mathcal{F}})=-\infty$.
- An isotrivial fibration of genus ≥ 2, if $\kappa(K_{\mathcal{F}})=1$.

This gives the "singular" analogous of the Proposition 3.3. Then, for instance, Hilbert modular surfaces which are minimal resolution of surfaces with cusps give examples of surfaces S with $c_1(S)^2<2\,c_2(S)$ such that $GG(S)=S$.

Finally, what if $K_{\mathcal{F}}$ is big? Here is a natural question which was asked to us by M. McQuillan in a private communication.

Question 3.4. *Let S be a surface admitting a holomorphic foliation by curve $\mathcal{F}$ with canonical singularities. Suppose that $K_{\mathcal{F}}$ is big. Is it then true that $GG(S,T_{\mathcal{F}})\subsetneq S$? If moreover S is of general type, what can be said about $GG(S)$ in this case?*

4. Corollary 1.4 for Demailly-Semple jets

In this section we would like to give a somehow more geometric proof of Corollary 1.4 in the case of invariant jet differentials, even if this case is of course included in what is proved above. For the sake of simplicity we shall assume that $L\subset T_X$ is a holomorphic line subbundle and not just an injection of sheaves. Invariant jet differentials were introduced in [Dem97] as a refined version of Green-Griffiths' ones: they are constructed taking invariant holomorphic functions on the k-jet space by a larger group, namely the full group of k-jets of biholomorphisms of $(\mathbb{C},0)$, instead of merely homotheties. The vector bundle $E_{k,m}V^*\to X$ of invariant jet differential of order k and weighted degree m acting on germs of holomorphic curves tangent to $V\subset T_X$ can be obtained as a direct image of an invertible sheaf on a tower of projective spaces, as follows.

Start with a directed manifold (X,V) and define the new directed manifold $(\widetilde{X},\widetilde{V})$ to be

$$\widetilde{X}=P(V),\quad \widetilde{V}=\pi_*^{-1}\big(\mathcal{O}_{P(V)}(-1)\big)\subset T_{P(V)},$$

where $\pi\colon P(V)\to X$ is the projectivized bundle of lines of V and $\mathcal{O}_{P(V)}(-1)\subset\pi^*V$ the tautological line bundle of $P(V)$. Now, set $(P_0(V),V_0)=(X,V)$ and define inductively

$$(P_k(V),V_k)=(\widetilde{P_{k-1}(V)},\widetilde{V_{k-1}})$$

together with the total projection $\pi_{0,k}\colon P_k(V)\to X$. The functoriality of this construction shows that if $(Y,W)\subset(X,V)$ is a directed submanifold (*i.e.* Y is a smooth submanifold of X, possibly the whole X, and $W\subset T_Y\subset T_X|_Y$ is a holomorphic subbundle of $V|_Y$), then for all positive integers k we have

$$P_k(W)\subset P_k(V),\quad W_k\subset V_k|_{P_k(W)}\quad\text{and}\quad \mathcal{O}_{P_k(V)}(-1)|_{P_k(W)}=\mathcal{O}_{P_k(W)}(-1),$$

and moreover the projection maps are of course compatible.

If $\varphi\colon(\mathbb{C},0)\to X$ is a germ of holomorphic curve tangent to V then we can define a projectivized lifting $\varphi_{[k]}\colon(\mathbb{C},0)\to P_k(V)$. Such a lifting is tangent to V_k and satisfies $\pi_{0,k}\circ\varphi_{[k]}=\varphi$.

It turns out that for any positive integer m there is an isomorphism of sheaves

$$(\pi_{0,k})_*\mathcal{O}_{P_k(V)}(m)\simeq\mathcal{O}_X(E_{k,m}V^*).$$

By Theorem 1.1, for any positive line bundle $A\to X$ and for all positive integers k,m, we have that

$$f_{[k]}(\mathbb{C})\subset\operatorname{Bs}\big(\mathcal{O}_{P_k(V)}(m)\otimes\pi_{0,k}^*A^{-1}\big),$$

whenever $f\colon \mathbb{C} \to X$ is an entire curve tangent to V (here, $\mathrm{Bs}(\bullet)$ stands for base locus of a line bundle).

With this is mind, we define the *Demailly-Semple locus* $DS(X, V)$ to be

$$DS(X, V) = DS_A(X, V),$$

where

$$DS_A(X, V) = \bigcap_{k\geq 1} \pi_{0,k}\left(\bigcap_{m\geq 1} \mathrm{Bs}\big(\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}\big)\right).$$

The fact that $DS_A(X, V)$ is independent of the ample line bundle $A \to X$ can be shown in the same way as in Lemma 2.2. The relation

$$\mathrm{Exc}(X, V) \subset GG(X, V) \subset DS(X, V)$$

is immediate from definitions and Theorem 1.1. *We do not know wether* $GG(X, V) = DS(X, V)$*, for instance, under the natural hypothesis that* $\det V^*$ *is big.*

Proof of Corollary 1.4 for Demailly-Semple jets. Fix any ample line bundle $A \to X$. We shall construct for each integer $k > 0$ a smooth manifold $X_k \subset P_k(V)$ which projects biholomorphically onto X via $\pi_{0,k}$ and such that for all integers $m > 0$ we have

$$X_k \subset \mathrm{Bs}\big(\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}\big).$$

The starting datum is a directed manifold (X, V) together with a rank one holomorphic subbundle $L \subset V \subset T_X$ such that $\kappa(L^{-1}) < \dim X$. We define X_k to be

$$X_k := P_k(L) \subset P_k(V).$$

Since at each step we are always projectivizing a rank one vector bundle, all X_k are isomorphic to the starting X, the isomorphism being given by the projections $\pi_{0,k}$. Moreover, since as we have seen

$$\mathcal{O}_{P_k(V)}(-1)|_{P_k(L)} = \mathcal{O}_{P_k(L)}(-1)$$

and, on the other hand,

$$\mathcal{O}_{P_k(L)}(-1) \simeq \pi_{0,k}^* L \simeq L,$$

we deduce that the restriction of $\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}$ to $X_k = P_k(L)$ has no nonzero holomorphic sections for all positive integers m by Kodaira's lemma, being isomorphic to $L^{-\otimes m} \otimes A^{-1}$. But then, it follows that the base locus of $\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}$ must necessarily contain X_k, and we are done. ☐

5. Final remarks on the Green-Griffiths-Lang conjecture

The final step for the solution of the Green-Griffiths-Lang conjecture using only jet bundles requires to show that at least one of the base loci

$$B_k := \bigcap_{m\geq 1} \mathrm{Bs}\big(\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}\big) \subset P_k(V)$$

projects down to a proper algebraic subvariety $Y_k := \pi_{0,k}(B_k)$ in X. These lines show that, unfortunately, this is hopeless without any further assumption on X besides that of being of general type.

Nevertheless, one could think of a less demanding property, namely that for all irreducible subvarieties $Z \subset P_k(V)$ such that $\pi_{0,k}(Z) = X$, the restriction $\mathcal{O}_{P_k(V)}(1)|_Z$ is big.

This is exactly what is proved in a theorem of Lu and Yau [LY90] for order one jets, for X a surface of general type with $c_1(X)^2 - 2\,c_2(X) > 0$, achieving thus the proof of the Green-Griffiths-Lang conjecture in this case. But we can not expect even this less demanding property to be true in general since we have seen in the previous section that there exist projective manifolds X of general type such that for each $k > 0$ there is a smooth submanifold $Z_k \subset P_k(V)$ which projects biholomorphically onto X via $\pi_{0,k}$ and such that for all integers $m > 0$ we have

$$X_k \subset \mathrm{Bs}\big(\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}\big).$$

Thus, we see that Lu and Yau's result is sharp and that $c_1^2(X) = 2\,c_2(X)$ is somehow a threshold for projective surfaces of general type as far as the solution to the Green-Griffiths-Lang conjecture using only base locus of sections of k-jet bundles is concerned.

On the other hand, McQuillan's celebrated work on surfaces [McQ98] shows that the conjecture is true whenever the second Segre number $c_1^2 - c_2$ of the surface is positive (this being the case for the compact free quotients of the bidisc thanks to the Hirzebruch proportionality principle, see [Hir58]). Observe that his proof relies upon a combination of the theories of jet differentials and that of foliations: jet differentials of order one (the existence of which is assured by the assumption on the Segre number) produce (multi)foliations such that every entire curve must be tangent to. The algebraic degeneracy is then obtained in this situation as a consequence of deep results about parabolic leaves of holomorphic foliations on surfaces of general type rather than trying to control the Green-Griffiths locus (which, *a posteriori* would not give the desired result, also in view of what is described in these lines).

References

[Bea00] Arnaud Beauville. Complex manifolds with split tangent bundle. In *Complex analysis and algebraic geometry*, pages 61–70. de Gruyter, Berlin, 2000.

[Bru97] Marco Brunella. Feuilletages holomorphes sur les surfaces complexes compactes. *Ann. Sci. École Norm. Sup. (4)*, 30(5):569–594, 1997.

[Bru04] Marco Brunella. *Birational geometry of foliations.* Publicações Matemáticas do IMPA. [IMPA Mathematical Publications]. Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2004.

[Dem97] Jean-Pierre Demailly. Algebraic criteria for Kobayashi hyperbolic projective varieties and jet differentials. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 285–360. Amer. Math. Soc., Providence, RI, 1997.

[Dem11] Jean-Pierre Demailly. Holomorphic Morse inequalities and the Green-Griffiths-Lang conjecture. *Pure Appl. Math. Q.*, 7(4, Special Issue: In memory of Eckart Viehweg):1165–1207, 2011.

[Fre90] Eberhard Freitag. *Hilbert modular forms.* Springer-Verlag, Berlin, 1990.

[GG80] Mark Green and Phillip Griffiths. Two applications of algebraic geometry to entire holomorphic mappings. In *The Chern Symposium 1979 (Proc. Internat. Sympos., Berkeley, Calif., 1979)*, pages 41–74. Springer, New York, 1980.

[Hir58] Friedrich Hirzebruch. Automorphe Formen und der Satz von Riemann-Roch. In *Symposium internacional de topología algebraica International symposium on algebraic topology*, pages 129–144. Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958.

[Kat92] Svetlana Katok. *Fuchsian groups.* Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1992.

[Lan86] Serge Lang. Hyperbolic and Diophantine analysis. *Bull. Amer. Math. Soc. (N.S.)*, 14(2):159–205, 1986.

[Lu96] Steven Shin-Yi Lu. On hyperbolicity and the Green-Griffiths conjecture for surfaces. In *Geometric complex analysis (Hayama, 1995)*, pages 401–408. World Sci. Publ., River Edge, NJ, 1996.

[LY90] Steven Shin-Yi Lu and S.-T. Yau. Holomorphic curves in surfaces of general type. *Proc. Nat. Acad. Sci. U.S.A.*, 87(1):80–82, 1990.

[McQ98] Michael McQuillan. Diophantine approximations and foliations. *Inst. Hautes Études Sci. Publ. Math.*, No. 87:121–174, 1998.

[McQ08] Michael McQuillan. Canonical models of foliations. *Pure Appl. Math. Q.*, 4(3, part 2):877–1012, 2008.

[McQ12] Michael McQuillan. McQuillan Theory F.A.Q., webpage available at the Url `http://www.mat.uniroma2.it/~mcquilla/faq/faq.html`, last update July 2012.

[Mer10] J. Merker. Complex projective hypersurfaces of general type: toward a conjecture of Green and Griffiths. *ArXiv e-prints*, May 2010.

[Mok89] Ngaiming Mok. *Metric rigidity theorems on Hermitian locally symmetric manifolds*, volume 6 of *Series in Pure Mathematics*. World Scientific Publishing Co. Inc., Teaneck, NJ, 1989.

[Rag72] M. S. Raghunathan. *Discrete subgroups of Lie groups*. Springer-Verlag, New York, 1972. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 68.

[SB95] N. I. Shepherd-Barron. Infinite generation for rings of symmetric tensors. *Math. Res. Lett.*, 2(2):125–128, 1995.

[Sha78] Ira H. Shavel. A class of algebraic surfaces of general type constructed from quaternion algebras. *Pacific J. Math.*, 76(1):221–245, 1978.

[Shi94] Goro Shimura. *Introduction to the arithmetic theory of automorphic functions*, volume 11 of *Publications of the Mathematical Society of Japan*. Princeton University Press, Princeton, NJ, 1994. Reprint of the 1971 original, Kanô Memorial Lectures, 1.

[SY97] Yum-Tong Siu and Sai-Kee Yeung. Defects for ample divisors of abelian varieties, Schwarz lemma, and hyperbolic hypersurfaces of low degrees. *Amer. J. Math.*, 119(5):1139–1172, 1997.

[Tak90] Kisao Takeuchi. Algebraic surfaces derived from unit groups of quaternion algebras. In *Number theory (Banff, AB, 1988)*, pages 529–552. de Gruyter, Berlin, 1990.

[Yau93] Shing-Tung Yau. A splitting theorem and an algebraic geometric characterization of locally Hermitian symmetric spaces. *Comm. Anal. Geom.*, 1(3-4):473–486, 1993.

Simone Diverio, CNRS et Institut de Mathématiques de Jussieu - Paris Rive Gauche, Université Pierre et Marie Curie, Paris.

E-mail address: `diverio@math.jussieu.fr`

Erwan Rousseau, Laboratoire d'Analyse, Topologie, Probabilités, Université d'Aix-Marseille et CNRS, Marseille.

E-mail address: `erwan.rousseau@cmi.univ-mrs.fr`